

References

1. Girard S, Tremblay L, Lepage M, Sébire G. IL-1 receptor antagonist protects against placental and neurodevelopmental defects induced by maternal inflammation. *J Immunol*. 2010 Apr 1; 184(7):3997-4005.
2. Hsiao EY, Patterson PH. Activation of the maternal immune system induces endocrine changes in the placenta via IL-6. *Brain Behav Immun*. 2011 May; 25(4):604-15.
3. Khan D, Fernando P, Cicvaric A, Berger A, Pollak A, Monje FJ, Pollak DD. Long-term effects of maternal immune activation on depression-like behavior in the mouse. *Transl Psychiatry*. 2014 Feb 18; 4: e363.
4. C. M. Wilke, K. Bishop, D. Fox, W. Zou, Deciphering the role of Th17 cells in human disease. *Trends Immunol*. 32, 603–611 (2011).
5. L. Y. Al-Ayadhi, G. A. Mostafa, Elevated serum levels of interleukin-17A in children with autism. *J. Neuroinflammation* 9, 158 (2012).
6. K. Suzuki, H. Matsuzaki, K. Iwata, Y. Kamenno, C. Shimmura, S. Kawai, Y. Yoshihara, T. Wakuda, K. Takebayashi, S. Takagai, K. Matsumoto, K. J. Tsuchiya, Y. Iwata, K. Nakamura, M. Tsujii, T. Sugiyama, N. Mori, Plasma cytokine profiles in subjects with high-functioning autism spectrum disorders. *PLOS ONE* 6, e20470 (2011).
7. Akintunde ME, Rose M, Krakowiak P, Heuer L, Ashwood P, Hansen R, Hertz-Picciotto I, Van de Water J. Increased production of IL-17 in children with autism spectrum disorders and co-morbid asthma. *J Neuroimmunol*. 2015 Sep 15; 286:33-41.
8. B. van der Zwaag, L. Franke, M. Poot, R. Hochstenbach, H. A. Spierenburg, J. A. Vorstman, E. van Daalen, M. V. de Jonge, N. E. Verbeek, E. H. Brilstra, R. van 't Slot, R. A. Ophoff, M. A. van Es, H. M. Blauw, J. H. Veldink, J. E. Buizer-Voskamp, F. A. Beemer, L. H. van den Berg, C. Wijmenga, H. K. van Amstel, H. van Engeland, J. P. Burbach, W. G. Staal, Gene-network analysis identifies susceptibility genes related to glycobiology in autism. *PLOS ONE* 4, e5324 (2009).
9. M. Mandal, A. C. Marzouk, R. Donnelly, N. M. Ponzio, Preferential development of Th17 cells in offspring of immunostimulated pregnant mice. *J. Reprod. Immunol*. 87, 97–100 (2010).
10. E. Y. Hsiao, S. W. McBride, J. Chow, S. K. Mazmanian, P. H. Patterson. Modeling an autism risk factor in mice leads to permanent immune dysregulation. *Proc. Natl. Acad. Sci. U.S.A.*
11. Ozkan ZS, Deveci D, Simsek M, Ilhan F, Risvanli A, Sapmaz E. What is the impact of SOCS3, IL-35 and IL17 in immune pathogenesis of recurrent pregnancy loss? *J Matern Fetal Neonatal Med*. 2015 Feb;28(3):324-8.

12. Zidan HE, Rezk NA, Alnemr AA, Moniem MI. Interleukin-17 and leptin genes polymorphisms and their levels in relation to recurrent pregnancy loss in Egyptian females. *Immunogenetics*. 2015 Oct 14.
13. Cornelius DC, Lamarca B. TH17- and IL-17- mediated autoantibodies and placental oxidative stress play a role in the pathophysiology of pre-eclampsia. *Minerva Ginecol*. 2014 Jun;66(3):243-9. Review.
14. Galazios G, Tsoulou S, Zografou C, Tripsianis G, Koutlaki N, Papazoglou D, Tsikouras P, Maltezos E, Liberis V. The role of cytokines IL-6 and IL-8 in the pathogenesis of spontaneous abortions. *J Matern Fetal Neonatal Med*. 2011 Oct;24(10):1283-5.
15. V. K. Kuchroo, A. Awasthi, Emerging new roles of Th17 cells. *Eur. J. Immunol*. 42, 2211–2214 (2012).
16. Udenze I, Amadi C, Awolola N, Makwe CC. The role of cytokines as inflammatory mediators in preeclampsia. *Pan Afr Med J*. 2015 Mar 10; 20:219.